

Subject: Intervertebral Stabilization Devices for Spinal Fusion		Original Effective Date: 9/18/19
Policy Number: MCP-343	Revision Date(s):	
Review Date: 9/16/20		
MCPC Approval Date: 9/18/19, 9/16/20		

DISCLAIMER

This Molina Clinical Policy (MCP) is intended to facilitate the Utilization Management process. It expresses Molina's determination as to whether certain services or supplies are medically necessary, experimental, investigational, or cosmetic for purposes of determining appropriateness of payment. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered (i.e., will be paid for by Molina) for a particular member. The member's benefit plan determines coverage. Each benefit plan defines which services are covered, which are excluded, and which are subject to dollar caps or other limits. Members and their providers will need to consult the member's benefit plan to determine if there are any exclusion(s) or other benefit limitations applicable to this service or supply. If there is a discrepancy between this policy and a member's plan of benefits, the benefits plan will govern. In addition, coverage may be mandated by applicable legal requirements of a State, the Federal government or CMS for Medicare and Medicaid members. CMS's Coverage Database can be found on the CMS website. The coverage directive(s) and criteria from an existing National Coverage Determination (NCD) or Local Coverage Determination (LCD) will supersede the contents of this Molina Clinical Policy (MCP) document and provide the directive for all Medicare members.¹

Contents

DISCLAIMER..... 1

Description of Procedure/Service/Pharmaceutical..... 1

Recommendation 2

Summary of Medical Evidence..... 2

Coding Information..... 3

References..... 3

DESCRIPTION OF PROCEDURE/SERVICE/PHARMACEUTICAL ^{2 18-20}

This MCP addresses the use of flexible intervertebral stabilization devices as an adjunct to spinal fusion procedures to provide immobilization and stabilization of spinal segments. These devices are designed to allow some degree of spinal flexibility following spinal fusion surgery.

The most common surgery for chronic nonspecific low back pain with lumbar disc degenerative changes is spinal fusion, a procedure that fuses two or more vertebral bodies together. The goal is to restrict spinal motion and remove the degenerated disc (the presumed cause of pain) in order to relieve symptoms. A variety of fusion techniques are used. All involve the placement of a bone graft between the vertebrae. Fusion can be performed with or without supplemental hardware (instrumentation), such as pedicle rods, plates, screws, or cages that function as an internal splint while the bone graft heals. Fusion alters the normal mechanics of the spine and is associated with an increase in long-term degenerative changes in adjacent spine segments. The standard spinal fusion procedure for rigid spinal fixation involves the use of pedicle screws, rods, cages and plates.

Intervertebral stabilization devices have been proposed as an alternative to the use of standard rigid frames to try to improve the disadvantages of rigid instrumentation and increase the outcome of spinal fusion surgery. These devices are fixed in place using pedicle screws which are attached to the vertebral bodies adjacent to the intervertebral space being fused but are designed using flexible materials which suggest stabilizing the joint while still providing some flexibility.

Examples of intervertebral stabilization devices that have currently been approved by the U.S. Food and Drug Administration (FDA) 510(k) clearance process as an adjunct to interbody fusion include but are not limited to the following: Isobar Spinal System (Alphatec Spine, Inc. Carlsbad, CA), Dynesys® System (Zimmer Inc., Minneapolis MN), BioFlex® (BioSpine Co., Ltd, Sungdong-gu, Seoul Korea) and the DSSTM Stabilization System (Paradigm Spine, LLC, New York, NY).

Note: Intervertebral Stabilization devices for spinal fusion in this MCP differ from Interspinous process fixation devices and Interspinous Decompression devices. Please see the following related MCP's for additional information concerning these devices: Interspinous Decompression Devices for Spinal Stenosis (X Stop, non-fusion Coflex) MCP-222 and Interspinous Process Fixation Devices for Spinal Fusion MCP-339.

RECOMMENDATION

Intervertebral Stabilization Devices as an adjunct to spinal fusion are considered experimental, investigational and/or unproven for any indication, due to insufficient clinical evidence of safety and efficacy in published peer-reviewed medical literature.

SUMMARY OF MEDICAL EVIDENCE³⁻¹⁷

Overall, there is a paucity of evidence in the peer-reviewed published medical literature to support the long-term safety and effectiveness of intervertebral stabilization devices when used in combination with interbody fusion or as a stand-alone procedure. The available studies are small, with short follow-up of two years so the long-term efficacy and safety of the procedure are not known. Large well designed randomized controlled trials are needed to demonstrate the clinical utility of intervertebral stabilization devices compared with established standard surgical approaches involving pedicle screw-rod-cage-plate fixation with lumbar fusion procedures. A summary of the relevant studies are outlined below.

Bieri et al, (2018) published an analysis of data from the International Spine Tango Registry on 202 individuals who used the DSS stabilization system and 269 individuals who underwent PLIF. At a mean follow-up of 3 years, there was not a statistically significant difference in the mean Core Outcomes Measure Index (COMI) score improvement (3.4 points in the DSS group and 3.2 points in the PLIF group), p=0.69. Matched pairs were

also similar in terms of back and leg pain relief, blood loss during surgery and complication rates. However, there were significantly fewer repeat surgeries after DSS (0.8 per 100 observed person-years) than with PLIF (2.9 per 100 observed person-years). The authors indicated that there are no published prospective comparative studies evaluating the DSS stabilization system. ³

Lee et al, (2016) conducted a meta-analysis of the literature on the efficacy of the Dynesys system to compare clinical and radiological outcomes between individuals who underwent surgery with Dynesys versus posterior lumbar interbody fusion (PLIF) for degenerative spinal disease. A total of 7 studies with 506 participants met the eligibility criteria. Only one was a randomized controlled trial (RCT), two were prospective cohort studies and four were retrospective cohort studies. Clinical and radiological outcomes, including the Oswestry Disability Index (ODI) and pain measured by a visual analogue scale (VAS), were assessed at baseline and again at two years. Pooled analyses did not find significant differences between the two surgical methods in change in the ODI or in back or leg pain VAS scores. Rates of complications and length of hospital stay were similar in the two groups. ⁸

Fu et al, (2014) evaluated the functional and radiological outcomes of dynamic stabilization in conjunction with spinal fusion in a prospective study of 36 participants with follow-up for 24 months who underwent posterior Isobar dynamic stabilization for single-level degenerative lumbar disc disease with instability (DLDI) and mild adjacent level degeneration. Outcomes were assessed using functional [visual analog scale (VAS) and Oswestry Disability Index (ODI)] and radiological data (resting, functional X-rays and MRI). At 24 months, functional outcomes demonstrated significant improvement in mean VAS score by 38.9 points (p<0.01) and ODI by 22.4 points (p<0.01). Results indicated that individuals with single-level DLDI and mild adjacent level degeneration treated with Isobar semi-rigid stabilization demonstrated improvement in functional scores 2 years postoperatively. However, disc degeneration at the adjacent and index levels appears to continue despite using semi-rigid dynamic stabilization. The authors stated that additional long-term follow-up is ongoing to provide more extensive information. ⁶

CODING INFORMATION THE CODES LISTED IN THIS POLICY ARE FOR REFERENCE PURPOSES ONLY. LISTING OF A SERVICE OR DEVICE CODE IN THIS POLICY DOES NOT IMPLY THAT THE SERVICE DESCRIBED BY THIS CODE IS COVERED OR NON-COVERED. COVERAGE IS DETERMINED BY THE BENEFIT DOCUMENT. THIS LIST OF CODES MAY NOT BE ALL INCLUSIVE.

CPT	Description
22899	Unlisted procedure, spine [when specified as insertion of a non-pedicle interspinous process fixation device]

REFERENCES

Government Agency

- Centers for Medicare & Medicaid Services (CMS). Medicare Coverage Database. Advanced Search: National Coverage Documents [search:]. Available at: <http://www.cms.gov/medicare-coverage-database/>
- Center for Devices and Radiological Health (CDRH). Premarket Approval (PMA) Database. Food and Drug Administration [website]. Available at: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMA/pma.cfm>

Peer Reviewed Publications

3. Bieri KS, Goodwin K, Aghayev E et al Dynamic posterior stabilization versus posterior lumbar intervertebral fusion: A matched cohort study based on the Spine Tango Registry. 2018; *J Neurol Surg*; 79:224-230.
4. Bothmann M, Kast E, Boldt GJ, Oberle J. Dynesys fixation for lumbar spine degeneration. *Neurosurg Rev*. 2007 Sep 29; [Epub ahead of print].
5. Bredin S, Demay O et al. Posterolateral fusion versus Dynesys dynamic stabilization: Retrospective study at a minimum 5.5years' follow-up. *Orthop Traumatol Surg Res*. 2017 Dec;103(8):1241-1244.
6. Fu L, France A, Xie Y, et al. Functional and radiological outcomes of semi-rigid dynamic lumbar stabilization adjacent to single-level fusion after 2 years. *Arch Orthop Trauma Surg*. 2014; 134(5):605-610.
7. Grob D, Benini A, Junge A, Mannion AF. Clinical experience with the Dynesys semirigid fixation system for the lumbar spine: surgical and patient-oriented outcome in 50 cases after an average of 2 years. *Spine*. 2005; 30(3):324-331.
8. Lee C-H, Jahng T-A, Hyan S-J et al. Dynamic stabilization using the Dynesys system versus posterior lumbar interbody fusion for the treatment of degenerative lumbar spinal disease; a clinical and radiological outcomes-based meta-analysis. *Neurosurg Focus* 2016; 50: 1-9.
9. Maida G, Altruda C, Gatti M, et al. Two-year follow-up after microsurgical discectomy and dynamic percutaneous stabilization in degenerate and herniated lumbar disc: clinical and neuroradiological outcome. *J Neurosurg Sci*. 2014; 58(2):95-102.
10. Pham M, Mehta V, Patel N, et.al. Complications associated with the Dynesys dynamic stabilization system: a comprehensive review of the literature. *Neurosurg Focus*. 2016; 40(1):E2.
11. Putzier M, Schneider SV, Funk JF, et al. The surgical treatment of the lumbar disc prolapse: nucleotomy with additional transpedicular dynamic stabilization versus nucleotomy alone. *Spine*. 2005; 30(5):E109-114.
12. Schaeren S, Broger I, Jeanneret B. Minimum four-year follow-up of spinal stenosis with degenerative spondylolisthesis treated with decompression and dynamic stabilization. *Spine*. 2008; 33(18):E636-642.
13. Schnake KJ, Schaeren S, Jeanneret B. Dynamic stabilization in addition to decompression for lumbar spinal stenosis with degenerative spondylolisthesis. *Spine*. 2006; 31(4):442-449.
14. Welch WC, Cheng BC, Awad TE, et al. Clinical outcomes of the Dynesys dynamic neutralization system: 1-year preliminary results. *Neurosurg Focus*. 2007 15; 22(1):E8.
15. Wu H, Pang Q, Jiang G. Medium-term effects of Dynesys dynamic stabilization versus posterior lumbar interbody fusion for treatment of multisegmental lumbar degenerative disease. *J Int Med Res* 2017; 45: 1562-1573.
16. Yang Y, Hong Y et al. Comparison of clinical and radiographic results between isobar posterior dynamic stabilization and posterior lumbar inter-body fusion for lumbar degenerative disease: A four-year retrospective study. *Clin Neurol Neurosurg*. 2015 Sep;136:100-6.
17. Zhang Y, Zhang ZC et al. Long-Term Outcome of Dynesys Dynamic Stabilization for Lumbar Spinal Stenosis. *Chin Med J (Engl)*. 2018 Nov 5;131(21):2537-2543.

Professional Society Guidelines

18. North American Spine Society (NASS). NASS Coverage Policy Recommendations. Interspinous fixation with fusion May 2014.
19. The American Association of Neurological Surgeons (AANS) Guideline update for the performance of fusion procedure for degenerative disease of the lumbar spine. Part 7: Lumbar fusion for intractable low-back pain without stenosis or spondylolisthesis. J Neurosurg Spine 21:42–47, 2014.

Other Resources

20. UpToDate: [website]. Waltham, MA: Walters Kluwer Health; 2020.
 - Chou R. Subacute and chronic low back pain: Surgical treatment.
 - Levin K. Lumbar spinal stenosis: Treatment and prognosis.
21. Advanced Medical Review (AMR): Policy reviewed by practicing MD board certified in Orthopaedic Surgery. 7/18/19

Review/Revision History:

9/18/19: New Policy

9/16/20: Policy reviewed, no changes. Added TOC.