

# Routine Screening and Lab Recommendations

	1 <sup>st</sup> Trimester (weeks 1-13)	2 <sup>nd</sup> Trimester (weeks 14-27)	3 <sup>rd</sup> Trimester (weeks 28-40+)	Postpartum
<b>Recommended checks at each prenatal visit</b>	Weight, Blood Pressure, Fetal Heart tones, Fetal position, Urine	Weight, Blood Pressure, Fetal Heart tones, Fetal position, Urine	Weight, Blood Pressure, Fetal Heart tones, Fetal position, Urine	Postpartum visit at 1-6 weeks depending on risk factors
<b>Recommended Labs</b>	Complete Blood Count (CBC) Blood Type (including Rh factor) Rubella Syphilis Hepatitis B (HBsAg) Hepatitis C HIV STI screening Urinalysis and Culture	Glucose Screening (24-28) Complete Blood Count for anemia screening	Hepatitis B (HBsAg) Hepatitis C Syphilis HIV Group B Strep Screening (36 weeks)	
<b>Recommended screenings</b>	Tobacco use screening Drug screening Depression Screening (PHQ9)	Depression Screening at 28 weeks	Depression Screening	Depression Screening
<b>Recommended Testing</b>	First trimester ultrasound for dating Option of prenatal genetic screening and carrier testing	Fetal Anatomy Ultrasound		
<b>Recommended Care</b>	Dental Visit (Member Incentive available)		Tobacco Use and Drug screening	Contraception plan

# Vaccination Recommendations in Pregnancy

Tdap	Influenza	RSV	Covid-19	Hepatitis A	Hepatitis B
<p>Weeks 27-36</p> <p>*Recommend family within household vaccinate as well.</p> <p>*Repeated every pregnancy.</p>	<p>Annually during Flu Season</p> <p>Recommended September-October but would recommend receiving anytime during flu season</p> <p>*Only inactivated or recombinant.</p> <p>*Recommend family within household vaccinate as well.</p>	<p>Weeks 32-36 (Sept-Jan)</p> <p>*Only Pfizer Abrysvo can be given during pregnancy</p> <p>*Only recommended during one pregnancy. If patient received in prior pregnancy, they would not receive immunization in subsequent pregnancies.</p>	<p>Any trimester of pregnancy.</p>	<p>Recommend during pregnancy if at high-risk (Illegal drug use, international travel, experiencing homelessness, international adoptee, HIV history, occupational risk).</p> <p>*offered as inactivated virus.</p>	<p>Recommend during pregnancy if not previously vaccinated.</p> <p>*Offered as non-infectious HBsAg virus.</p> <p>*Energix-B, Recombivax HB, Twinrix</p>