



Original Effective Date: 11/01/2016
Current Effective Date: 03/03/2023
Last P&T Approval/Version: 01/25/2023
Next Review Due By: 01/2024
Policy Number: C9837-A

Pulmonary Arterial Hypertension (PAH)

PRODUCTS AFFECTED

Endothelin-Receptor Antagonists (ERAs)

Letairis (ambrisentan), Tracleer (bosentan), Opsumit (macitentan), ambrisentan, bosentan

Phosphodiesterase type 5 inhibitors (PDE-5 inhibitors)

Adcirca (tadalafil), ALYQ (tadalafil), Tadalafil (tadalafil), Revatio (sildenafil), sildenafil, tadalafil

Soluble Guanylate Cyclase Stimulator

Adempas (riociguat)

Prostanoids/prostacyclin therapies

Uptravi (selexipag), Epoprostenol, Orenitram (treprostinil extended-release tablets), Tyvaso (treprostinil inhalation solution), Veletri (epoprostenol for injection), Remodulin (treprostinil injection), Ventavis (iloprost), Flolan (epoprostenol for injection)

COVERAGE POLICY

Coverage for services, procedures, medical devices and drugs are dependent upon benefit eligibility as outlined in the member's specific benefit plan. This Coverage Guideline must be read in its entirety to determine coverage eligibility, if any.

This Coverage Guideline provides information related to coverage determinations only and does not imply that a service or treatment is clinically appropriate or inappropriate. The provider and the member are responsible for all decisions regarding the appropriateness of care. Providers should provide Molina Healthcare complete medical rationale when requesting any exceptions to these guidelines.

Documentation Requirements:

Molina Healthcare reserves the right to require that additional documentation be made available as part of its coverage determination; quality improvement; and fraud; waste and abuse prevention processes. Documentation required may include, but is not limited to, patient records, test results and credentials of the provider ordering or performing a drug or service. Molina Healthcare may deny reimbursement or take additional appropriate action if the documentation provided does not support the initial determination that the drugs or services were medically necessary, not investigational or experimental, and otherwise within the scope of benefits afforded to the member, and/or the documentation demonstrates a pattern of billing or other practice that is inappropriate or excessive.

DIAGNOSIS:

Pulmonary Arterial Hypertension, Chronic Thromboembolic Pulmonary Hypertension, Pulmonary hypertension associated with interstitial lung disease

REQUIRED MEDICAL INFORMATION:

This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. If a drug within this policy receives an updated FDA label within the last 180 days,

Molina Healthcare, Inc. confidential and proprietary © 2023

This document contains confidential and proprietary information of Molina Healthcare and cannot be reproduced, distributed, or printed without written permission from Molina Healthcare. This page contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with Molina Healthcare.

Drug and Biologic Coverage Criteria

medical necessity for the member will be reviewed using the updated FDA label information along with state and federal requirements, benefit being administered and formulary preferencing. Coverage will be *determined on a case-by case basis until the criteria can be updated through Molina Healthcare, Inc. clinical governance. Additional information may be required on a case-by-case basis to allow for adequate review.*

FOR Phosphodiesterase type 5 inhibitors (PDE-5 inhibitors) Adcirca (tadalafil), ALYQ (tadalafil), Tadalafil (tadalafil), Revatio (sildenafil), sildenafil, tadalafil ONLY:

FOR NEW YORK HEALTHPLAN MEMBERS ONLY: Reviewer MUST check the Erectile Dysfunction Verification System (EDVS) for each request to determine member's sex offender status. IF a member is on the sex offender list the request must be forwarded to the medical director AND provider must provide the rationale for prescribing a PDE-5 inhibitors and note the reason(s) why alternative treatment options are inappropriate to treat the enrollee's health condition. Before issuing an adverse determination for a prescribed PDE5 inhibitor, the Medical Director must make reasonable attempts to engage in a peer-to-peer discussion with the requesting provider to understand the reasons behind the need for prescribing the requested PDE5 inhibitor or drug. The Medical Director may extend the review time, if requested by the provider or patient, or if such extension is in the best interest of the patient's health condition.

For after-hours, holiday, and weekend pharmacy requests for prescription PDE5 inhibitors Molina Healthcare, Inc can authorize a seventy-two (72) hour emergency supply of the prescription PDE5 inhibitor and must note within the authorization file that the drug is prescribed to treat a condition other than sexual or erectile dysfunction and that the drug has been approved by the FDA to treat that condition. The case must still be sent to the Medical director for checking the EDVS status of the member, as soon as possible on the next business day.

A. PULMONARY ARTERIAL HYPERTENSION:

1. Documented diagnosis of pulmonary arterial hypertension (PAH)
AND
2. Documentation of member's WHO group and WHO functional class
AND
3. Medication requested for treatment is consistent with its own FDA- labeled WHO functional class (see Appendix)
AND
4. FOR ORENITAM (TREPROSTINIL) and UPTRAVI (SELEXIPAG) ONLY:
(a) Member has tried or is currently receiving two oral therapies for PAH from two of the three following different categories (either alone or in combination) each for ≥ 60 days: one phosphodiesterase type 5 (PDE5) inhibitor, one endothelin receptor antagonist (ERA), OR Adempas (riociguat)
OR
(b) Member is receiving or has received in the past for PAH one prostacyclin therapy, or a prostacyclin receptor agonist for PAH.
AND
5. FOR ADCIRCA, ALYQ, TADLIQ, REVATIO, TADALAFIL OR SILDENAFIL ONLY: The member will not be taking another PDE5 inhibitor at the same time as the requested therapy
AND
6. FOR REQUESTS FOR OPSUMIT (MACITENTAN): Documentation of a trial and failure or contraindication to ambrisentan and bosentan
AND
7. IF REQUEST IS FOR BRAND PRODUCT WITH GENERIC AVAILABLE: Documentation the patient has failed a trial of the respective generic product and/or the patient cannot take the respective generic product due to a formulation difference in the active ingredient or due to a formulation difference in the inactive ingredient(s) [e.g., difference in dyes, fillers,

Drug and Biologic Coverage Criteria

preservatives between the brand and the generic product which would result in a significant allergy or serious adverse reaction per the prescribing physician [DOCUMENTATION REQUIRED].

B. CHRONIC THROMBOEMBOLIC PULMONARY HYPERTENSION (CTEPH)- ADEMPAS ONLY:

1. Documented diagnosis of chronic thromboembolic pulmonary hypertension (CTEPH)
AND
2. (a) Documentation member has been diagnosed as inoperable by a center specializing in CTEPH or pulmonary endarterectomy (PEA)
OR
(b) Documentation member been diagnosed as recurrent or persistent CTEPH after PEA
AND
3. Prescriber attests Adempas (riociguat) will NOT be administered with phosphodiesterase (PDE)-5 inhibitors or nitrates

C. PULMONARY HYPERTENSION ASSOCIATED WITH INTERSTITIAL LUNG DISEASE-TYVASO ONLY

1. Documentation member has a diagnosis of an interstitial lung disease [examples: Idiopathic pulmonary fibrosis (IPF) or pulmonary fibrosis (PF); Combined pulmonary fibrosis and emphysema (CPFE); Connective tissue disease (CTD)]
AND
2. Documentation that member has Group 3 pulmonary hypertension
AND
3. Documentation of member's baseline 6-minute walk test [DOCUMENTATION REQUIRED]

CONTINUATION OF THERAPY:

A. PULMONARY ARTERIAL HYPERTENSION:

1. Documentation that the member has demonstrated a beneficial response to therapy per the prescribing physician (e.g., Improvement in Six Minute Walking Test or Exercise Capacity, Improvement in WHO Functional Class, decrease in mean pulmonary artery pressure [mPAP], Increase in Cardiac Index, etc.)
AND
2. Prescriber attests to or clinical reviewer has found no evidence of intolerable adverse effects or drug toxicity

B. CHRONIC THROMBOEMBOLIC PULMONARY HYPERTENSION (CTEPH) - ADEMPAS ONLY:

1. Documentation that member's condition has not progressed or worsened on therapy
AND
2. Prescriber attests to or clinical reviewer has found no evidence of intolerable adverse effects or drug toxicity

C. PULMONARY HYPERTENSION ASSOCIATED WITH INTERSTITIAL LUNG DISEASE-TYVASO ONLY

1. Documentation that member has had improvement in 6-minute walk test from baseline [DOCUMENTATION REQUIRED]
AND
2. Prescriber attests to or clinical reviewer has found no evidence of intolerable adverse effects or drug toxicity

DURATION OF APPROVAL:

Initial authorization: 6 months, Continuation of therapy: 12 months

NEW YORK PATIENTS ONLY**PLEASE NOTE: FOR ANY MEMBER ON THE SEX OFFENDER LIST, APPROVAL CAN ONLY BEFOR 30 DAYS PER AUTHORIZATION******

Drug and Biologic Coverage Criteria

PRESCRIBER REQUIREMENTS:

Prescribed by a board-certified cardiologist, pulmonologist, or physician affiliated with a center of expertise in pulmonary arterial hypertension.

AGE RESTRICTIONS:

No requirement

QUANTITY:

Adcirca (tadalafil), Alyq (tadalafil) 20 mg tablet 60 tabs per 30 days
Adempas (riociguat) 0.5 mg, 1mg, 1.5mg, 2mg, or 2.5mg tablet 90 tabs per 30 days
Letairis (ambrisentan) 5 mg or 10mg tablet 30 tabs per 30 days
Opsumit (macitentan) 10 mg tablet 30 tabs per 30 days
Orenitram (treprostinil): determined by tolerability, 90 tabs per 30 days
Revatio (sildenafil): 20 mg tablet 3 tablets per day OR 10 mg/mL oral suspension 2 bottles (224 mL)/30 days
Tracleer (bosentan) 62.5 mg or 125mg tablet 60 tabs per 30 days
Tyvaso (inhaled treprostinil) 0.6 mg/mL System Starter Kit 1 kit/180 days or 0.6 mg/mL System Refill kit -1 package of 28 ampules/28 days or 4 pack Carton 7 packages of 4 ampules/28 days
Uptravi (selexipag) tabs Titration pack 1 pack/180 days 200 mcg, 400 mcg, 600 mcg, 800 mcg, 1000 mcg, 1200 mcg, 1400 mcg or 1600 mcg tablet - 2 tablets per day
Ventavis (iloprost) 10 mcg/mL or 20 mcg/mL - 9 packages of 30 ampules/30 days
NOTE: For Uptravi and Orenitram: authorizations may occur in small quantities per strength to allow for titration as prescribed

PLACE OF ADMINISTRATION:

The recommendation is that oral medications in this policy will be for pharmacy benefit coverage and patient self-administered.

The recommendation is that inhalation medications in this policy will be for pharmacy or medical benefit coverage administered in a place of service that is a non-hospital facility-based location as per the Molina Health Care Site of Care program.

The recommendation is that infused medications in this policy will be for pharmacy or medical benefit coverage administered in a place of service that is a non-hospital facility-based location as per the Molina Health Care Site of Care program.

Note: Site of Care Utilization Management Policy applies for epoprostenol (generic for Flolan), Flolan (epoprostenol), Remodulin (treprostinil sodium), Tyvaso (treprostinil). For information on site of care, see [Specialty Medication Administration Site of Care Coverage Criteria \(molinamarketplace.com\)](https://www.molinamarketplace.com/specialty-medication-administration-site-of-care-coverage-criteria)

DRUG INFORMATION

ROUTE OF ADMINISTRATION:

Oral, Intravenous, Inhalation

DRUG CLASS:

Pulmonary Hypertension - Prostacyclin Receptor Agonist, Pulm Hyperten-Soluble Guanylate Cyclase Stimulator (sGC), Pulmonary Hypertension - Phosphodiesterase Inhibitors, Pulmonary Hypertension - Endothelin Receptor Antagonists, Prostaglandin Vasodilators

FDA-APPROVED USES:

Adcirca, Alyq, Tadliq (tadalafil): indicated for the treatment of pulmonary arterial hypertension (PAH) (WHO Group 1) to improve exercise ability.

Drug and Biologic Coverage Criteria

Adempas (riociguat): indicated for:

- Persistent/recurrent Chronic Thromboembolic Pulmonary Hypertension (CTEPH) (WHO Group 4) after surgical treatment or inoperable CTEPH to improve exercise capacity and WHO functional class
- Pulmonary Arterial Hypertension (PAH) (WHO Group 1) to improve exercise capacity, improve WHO functional class and to delay clinical worsening.

Flolan, Veletri (epoprostenol): indicated for the treatment of pulmonary arterial hypertension (PAH) (WHO Group 1) to improve exercise capacity.

Letairis (ambrisentan): indicated for the treatment of adults with pulmonary arterial hypertension World Health Organization [WHO] Group 1.

Remodulin (treprostinil injection): indicated for treatment of pulmonary arterial hypertension (PAH; WHO Group 1) to diminish symptoms associated with exercise and for patients who require transition from epoprostenol, to reduce the rate of clinical deterioration.

Revatio (sildenafil): indicated for the treatment of pulmonary arterial hypertension (PAH) (WHO Group I) in adults to improve exercise ability and delay clinical worsening.

Opsumit (macitentan): indicated for the treatment of pulmonary arterial hypertension (PAH, WHO Group I) to reduce the risks of disease progression and hospitalization for PAH

Orenitram (treprostinil extended-release tablets): indicated for the treatment of pulmonary arterial hypertension (PAH) (WHO Group 1) to delay disease progression and to improve exercise capacity.

Tracleer (bosentan): indicated for the treatment of pulmonary arterial hypertension (PAH) (WHO Group 1) in:

- adults to improve exercise ability and to decrease clinical worsening. Studies establishing effectiveness included predominantly patients with WHO Functional Class II-IV symptoms and etiologies of idiopathic or heritable PAH (60%), PAH associated with connective tissue diseases (21%), and PAH associated with congenital heart disease with left-to-right shunts (18%).
- pediatric patients aged 3 years and older with idiopathic or congenital PAH to improve pulmonary vascular resistance (PVR), which is expected to result in an improvement in exercise ability.

Tyvaso (treprostinil inhalation solution): indicated for:

- Pulmonary arterial hypertension (PAH; WHO Group 1) to improve exercise ability. Studies establishing effectiveness predominately included patients with NYHA Functional Class III symptoms and etiologies of idiopathic or heritable PAH (56%) or PAH associated with connective tissue diseases (33%).
- Pulmonary hypertension associated with interstitial lung disease (PH-ILD; WHO Group 3) to improve exercise ability.

Upravi (selexipag): indicated for the treatment of pulmonary arterial hypertension (PAH) World Health Organization (WHO) Group 1 to delay disease progression and reduce the risk of hospitalization for PAH.

Ventavis (iloprost): indicated for the treatment of PAH (World Health Organization [WHO] Group 1) to improve exercise tolerance, symptoms (NYHA Class), and lack of deterioration

COMPENDIAL APPROVED OFF-LABELED USES:

None


APPENDIX

APPENDIX:

Molina Healthcare, Inc. confidential and proprietary © 2023

This document contains confidential and proprietary information of Molina Healthcare and cannot be reproduced, distributed, or printed without written permission from Molina Healthcare. This page contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with Molina Healthcare.

Drug and Biologic Coverage Criteria



Treatment algorithm. PAH: pulmonary arterial hypertension; IPAH: idiopathic PAH; HPAH: heritable PAH; DPAH: drug-induced PAH; CCB: calcium channel blocker; PCA: prostacyclin analogue; PH: pulmonary hypertension. ^a: 2015 ESC/ERS PH guidelines Table 16; ^b: 2015 ESC/ERS PH guidelines Table 17; ^c: 2015 ESC/ERS PH guidelines Table 18; ^d: 2015 ESC/ERS PH guidelines Table 13; ^e: 2015 ESC/ERS PH guidelines Table 19; ^f: 2015 ESC/ERS PH guidelines Table 20; ^g: 2015 ESC/ERS PH guidelines Table 14; ^h: 2015 ESC/ERS PH guidelines Table 21; ⁱ: maximal medical therapy is considered triple combination therapy including a s.c. or an i.v. PCA (i.v. preferred in high-risk status); ^j: 2015 ESC/ERS PH guidelines Table 22

Risk stratification and medical therapy of pulmonary arterial hypertension

Nazzareno Galiè, Richard N. Channick, Robert P. Frantz, Ekkehard Grünig, Zhi Cheng Jing, Olga Moiseeva, Ioana R. Preston, Tomas Pulido, Zeenat Safdar, Yuichi Tamura, Vallerie V. McLaughlin *European Respiratory Journal* 2019 53: 1801889; DOI: 10.1183/13993003.01889-2018

Brand (generic)	WHO Group	WHO Functional Class
Adcirca (tadalafil)	Group 1 PAH	Functional Class II-III
Adempas (riociguat)	Group 1 PAH	Functional Class II-III
	Group 4 CTEPH	n/a
Alyq (tadalafil)	Group 1 PAH	Functional Class II-III
Flolan (epoprostenol)	Group 1 PAH	Functional Class III-IV

Molina Healthcare, Inc. confidential and proprietary © 2023

This document contains confidential and proprietary information of Molina Healthcare and cannot be reproduced, distributed, or printed without written permission from Molina Healthcare. This page contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with Molina Healthcare.

Drug and Biologic Coverage Criteria

sodium)		
Letairis (ambrisentan)	Group 1 PAH	Functional Class II-III
Opsumit (macitentan)	Group 1 PAH	Functional Class II-III
Orenitram (treprostinil)	Group 1 PAH	Functional Class II-III
Remodulin (treprostinil)	Group 1 PAH	Functional Class II-IV
Revatio (sildenafil)	Group 1 PAH	Functional Class II-III
Tracleer (bosentan)	Group 1 PAH	Functional Class II-IV
Tyvaso (treprostinil)	Group 1 PAH	Functional Class III
	Group 3 PH-ILD	n/a
Uptravi (selexipag)	Group 1 PAH	Functional Class II-III
Veletri (epoprostenol)	Group 1 PAH	Functional Class II-IV
Ventavis (iloprost)	Group 1 PAH	Functional Class III-IV

BACKGROUND AND OTHER CONSIDERATIONS

BACKGROUND:

Adempas, a soluble guanylate cyclase (sGC) stimulator, is indicated for the treatment of adults with persistent/recurrent chronic thromboembolic pulmonary hypertension (CTEPH) [World Health Organization {WHO} Group 4] after surgical treatment, or inoperable CTEPH, to improve exercise capacity and WHO functional class. *Adempas* is also indicated for the treatment of adults with pulmonary arterial hypertension (PAH) [WHO Group 1], to improve exercise capacity, WHO functional class and to delay clinical worsening. Efficacy in WHO Group 1 PAH was established in patients receiving *Adempas* as monotherapy or in combination with endothelin receptor antagonists (ERAs) or prostanoids. Studies establishing effectiveness included mainly patients with WHO functional class II or III and etiologies of idiopathic or heritable PAH (61%) or PAH associated with connective tissue diseases (25%). The starting dose of *Adempas* is 1 mg three times daily (TID) and can be titrated to 2.5 mg TID.

Risk Evaluation and Mitigation Strategies (REMS) Program Because of the risk of embryo-fetal toxicity associated with *Adempas* therapy, *Adempas* are available through a restricted program under the REMS. Under the REMS, only certified healthcare providers and pharmacies may prescribe and distribute *Adempas*.

Tracleer, *Letairis* and *Opsumit* are oral endothelin receptor antagonists (ERAs) that are used for the treatment of pulmonary arterial hypertension (PAH). *Tracleer*, which is given twice daily (BID), is indicated for the treatment of PAH (World Health Organization [WHO] Group 1) to improve exercise ability and decrease the rate of clinical worsening. Studies establishing the effectiveness included predominantly those with New York Heart Association (NYHA) Functional Class II to IV symptoms and etiologies of idiopathic or heritable PAH (60%), PAH associated with connective tissue diseases (21%), and PAH associated with congenital systemic-to-pulmonary shunts (18%). Patients with WHO Class II symptoms demonstrated reduction in the rate of clinical deterioration and a trend for improvement in walk distance. Physicians should consider if these benefits are sufficient to offset the risk of liver injury in WHO Class II patients, which may preclude future use as disease progression occurs.¹ *Letairis*, which is given once daily (QD), is indicated for the treatment of PAH (WHO Group 1) to improve exercise ability and delay clinical worsening; it is also indicated for use in combination with *Adcirca*® (tadalafil tablets) to reduce

Drug and Biologic Coverage Criteria

the risks of disease progression and hospitalization for worsening PAH, and to improve exercise ability.² Studies establishing effectiveness included predominantly those with WHO Functional Class II to III symptoms and etiologies of idiopathic or heritable PAH (60%) or PAH associated with connective tissue diseases (34%).² Opsumit, which is given QD, is indicated for the treatment of PAH (WHO Group 1) to delay disease progression.³ Disease progression included: death, initiation of intravenous or subcutaneous prostanoids, or clinical worsening of PAH (decreased 6-minute walk distance, worsening PAH symptoms, and need for additional PAH treatment). Opsumit also reduced hospitalizations for PAH. All agents are in Pregnancy Category X and have a Boxed Warning regarding teratogenicity. 1-3 Tracleer has a Boxed Warning regarding hepatotoxicity.¹ All agents have a Boxed Warning regarding embryofetal toxicity

Risk Evaluation and Mitigation Strategies (REMS) Program Because of the risk of embryo-fetal toxicity and hepatotoxicity (Tracleer) associated with Tracleer, Letairis, Opsumit therapy, Tracleer, Letairis, Opsumit are available through a restricted program under the REMS. Under the REMS, only certified healthcare providers and pharmacies may prescribe and distribute Tracleer, Letairis, Opsumit.

Epoprostenol injection is a prostacyclin vasodilator. It is indicated for the treatment of pulmonary arterial hypertension (PAH) World Health Organization (WHO) Group 1 to improve exercise capacity. Studies establishing the effectiveness predominately included patients with New York Heart Association (NYHA) Functional Class III to IV symptoms and etiologies of idiopathic or heritable PAH or PAH associated with connective tissue diseases. Several studies have noted beneficial effects with epoprostenol therapy. Epoprostenol is given by intravenous infusion through a central venous catheter.

Ventavis and Tyvaso are both inhaled prostacyclin vasodilators indicated for the treatment of pulmonary arterial hypertension (PAH). Ventavis, which is given six to nine times per day, is indicated for the treatment of PAH (World Health Organization [WHO] Group 1) to improve a composite endpoint consisting of exercise tolerance, symptoms (based on New York Heart Association [NYHA] Class), and lack of deterioration. Studies establishing effectiveness involved mainly patients with NYHA Functional Class III to IV symptoms and etiologies of idiopathic or heritable PAH (65%) or PAH associated with connective tissue diseases (23%). Tyvaso, which is given four times per day, is indicated for the treatment of PAH (WHO Group 1) to improve exerciseability.² Studies establishing effectiveness mainly included those with NYHA Functional Class III symptoms and etiologies of idiopathic or heritable PAH (56%) or PAH associated with connective tissue diseases (33%). An updated treatment algorithm (2013) by the 2nd World Symposium on Pulmonary Hypertension (WSPH) states that patients with Functional Class II should be treated initially with oral therapies (e.g., Adempas® [riociguat tablets], sildenafil [Revatio®], generics {Note: brand name Revatio injection also available}, Adcirca® [tadalafil tablets], Opsumit® [macitentan tablets], Tracleer® [bosentan tablets], and Letairis® [ambrisentan tablets]).⁷ Ventavis and Tyvaso are recommended for patients in Functional Class III and IV. In situations of inadequate response, combination therapy (including double or triple therapy) is recommended.

Orenitram, an oral prostacyclin vasodilator, is indicated for the treatment of pulmonary arterial hypertension (PAH) World Health Organization (WHO) Group 1 to improve exercise capacity. The trial that established the efficacy of Orenitram included mainly patients with WHO functional class II to III symptoms. The prescribing information notes that Orenitram is probably most useful to replace subcutaneous, intravenous, or inhaled treprostinil, but this use requires further study. The recommended starting dose is 0.25 mg twice daily (BID) with food, taken approximately 12 hours apart. Dosing is individualized and titrated to response and tolerability; increase in increments of 0.25 to 0.5 mg BID every 3 to 4 days.

Revatio and Adcirca are phosphodiesterase type 5 (PDE5) inhibitors indicated for the treatment of pulmonary arterial hypertension (PAH). Revatio is indicated for PAH (World Health Organization [WHO] Group I) in adults to improve exercise ability and delay clinical worsening. The delay in clinical worsening was demonstrated when Revatio was added to background epoprostenol injection therapy (Flolan® [generic], Veletri®). Studies establishing its effectiveness were short-term (12 to 16 weeks) and included

Drug and Biologic Coverage Criteria

mainly patients with New York Heart Association (NYHA) Functional Class II to III symptoms and idiopathic etiology (71%) or associated with connective tissue disease (25%). A limitation of use is that adding Revatio to Tracleer® (bosentan tablets) does not result in any beneficial impact on exercise capacity. The recommended dose of Revatio is 5 mg or 20 mg three times daily (TID) given approximately 4 to 6 hours apart. In the clinical trial no greater efficacy was achieved with the use of higher doses. Treatment with doses higher than 20 mg TID is not recommended. Revatio has a Warning regarding mortality with increasing doses in pediatric patients. In a long-term trial involving pediatric patients with PAH, an increase in mortality with increasing Revatio dose was noted. Deaths were first observed following about 1 year and causes of death were usual of those with PAH. Revatio, especially chronic use, is not recommended in children.¹ Adcirca is indicated for the treatment of PAH (WHO Group I) to improve exercise ability.² Studies establishing effectiveness were mainly in patients with NYHA Functional Class II to III symptoms and etiologies of idiopathic or heritable PAH (61%) or PAH associated with connective tissue diseases (23%). The recommended dose is 40 mg once daily (QD). Dividing the dose (40 mg) over the course of the day is not recommended.

Remodulin is a prostacyclin vasodilator indicated for the treatment of pulmonary arterial hypertension (PAH) [World Health Organization {WHO} Group 1] to diminish symptoms associated with exercise. Studies establishing the effectiveness involved those with New York Heart Association (NYHA) Functional Class II to IV symptoms and etiologies of idiopathic or heritable PAH (58%), PAH associated with congenital systemic-to-pulmonary shunts (23%), or PAH associated with connective tissue diseases (19%). Remodulin may be administered via continuous subcutaneous (SC) infusion or continuous intravenous infusion. However, due to the risks associated with chronic indwelling central venous catheters, including serious blood stream infections, continuous intravenous infusion should be reserved for those intolerant of the SC, or in whom these risks are still considered acceptable. In those with PAH requiring transition from epoprostenol injection, Remodulin is indicated to diminish the rate of clinical deterioration. The risks and benefits of each agent should be considered carefully before transition.¹ Several trials have shown benefits of Remodulin therapy.

Uptravi (selexipag), an oral prostacyclin receptor agonist, is indicated for the treatment of pulmonary arterial hypertension (PAH) World Health Organization (WHO) Group 1 to delay disease progression and reduce the risk of hospitalization. The trial that established the efficacy of selexipag included mainly patients with WHO functional class II to III symptoms. The recommended starting dose is 200 mcg twice daily (BID) and may be more tolerable when taken with food. Dosing is individualized and titrated to tolerability; increased in increments of 200 mcg weekly to up to 1600 mcg twice daily. Those with moderated hepatic impairment should begin with 200 mcg and only dose once daily. Selexipag is not recommended to be used with those that have severe hepatic impairment. Selexipag is not to be chewed, crushed, or split.

CONTRAINDICATIONS/EXCLUSIONS/DISCONTINUATION:

All other uses of pulmonary arterial hypertension therapies are considered experimental/investigational and therefore will follow the Molina Healthcare, Inc. off-label policy.

Contraindications to Adcirca, ALYQ, Tadalafil, Tadalafil include: concomitant use of organic nitrates or guanylate cyclase (GC) stimulators, history of known serious hypersensitivity reaction to tadalafil or Cialis.

Contraindications to Adempas (riociguat) include: pregnancy, use with nitrates or nitric oxide donors in any form, use with PDE inhibitors, patients with concomitant use of other soluble guanylate cyclase (sGC) stimulators, and pulmonary hypertension associated with idiopathic interstitial pneumonias (PH-IIP).

Contraindications to Letairis (ambrisentan): include pregnancy and idiopathic pulmonary fibrosis.

Contraindications to Tracleer (bosentan) include: pregnancy, use with cyclosporine, use with glyburide, and hypersensitivity.

Contraindications to Opsumit (macitentan) include: pregnancy, and hypersensitivity.

Contraindications to Orenitram (treprostinil) include: severe hepatic impairment (Child Pugh Class C).

Drug and Biologic Coverage Criteria

Contraindications to Tyvaso (treprostinil), Tyvaso DPI (treprostinil), and Remodulin (treprostinil) include:

No labeled contraindications.

Contraindications to Flolan (epoprostenol sodium) include: heart failure with reduced ejection fraction and hypersensitivity to Flolan and any of its ingredients.

Contraindications to Veletri (epoprostenol) include: congestive heart failure due to severe left ventricular systolic dysfunction, pulmonary edema, hypersensitivity to the drug or to structurally related compounds.

Contraindications to Revatio (sildenafil) include use with organic nitrates or riociguat and history of hypersensitivity reaction to sildenafil or any component of the tablet, injection or oral suspension.

Contraindications to Uptravi (selexipag) include: concomitant use with strong CYP2C8 inhibitors, hypersensitivity to the active substance or to any of the excipients.

Contraindications to Ventavis (iloprost) include: No labeled contraindications.

OTHER SPECIAL CONSIDERATIONS:

Treatment Guidelines The 6th World Symposium on Pulmonary Hypertension evidence-based treatment algorithm:

Initial approach

- After confirmation of the diagnosis of the treatment-naïve PAH patient in an expert centre, the suggested initial approach is the adoption of *general measures* and the initiation of *supportive therapy* (2015 ESC/ERS PH guidelines).
- *Acute vasoreactivity testing* should be performed to predict response to calcium channel blocker (CCBs) only in patients with IPAH, HPAH, and PAH associated with drugs and toxin use. Vasoreactive patients (see the Task Force article by Simonneau *et al.* in this issue of the *European Respiratory Journal*) should be treated with *high doses (progressively titrated) of CCBs*; adequate response should be confirmed after 3–6 months of treatment (2015 ESC/ERS PH guidelines). *Adequate treatment response to high doses of CCBs* is considered WHO FC I/II with sustained haemodynamic improvement (same or better than achieved in the acute test) after at least 1 year on CCBs only. Vasoreactive patients without an adequate treatment response to high doses of CCBs should be treated with approved PAH medications according to the non-vasoreactive patients' treatment strategy.
- *Non-responders to acute vasoreactivity testing who are at low or intermediate risk* should be treated with *initial oral combination therapy with an ERA and a PDE5i* (2015 ESC/ERS PH guidelines).
- Some specific PAH subsets in which the efficacy/safety ratio of initial combination therapy is not established should be treated with initial monotherapy.

Recommendations for initial monotherapy are reported in the 2015 ESC/ERS PH guidelines

- If initial monotherapy is chosen, as head-to-head comparisons among different compounds are not available, no evidence-based first-line monotherapy can be proposed. The choice of drug may depend on a variety of factors, including approval status, labeling, route of administration, side-effect profile, potential interaction with background therapies, patient preferences, comorbidities, physician experience and cost.
- *In non-vasoreactive and treatment-naïve patients at high risk*, initial combination therapy including *i.e.*, PCAs is recommended (2015 ESC/ERS PH guidelines). Intravenous epoprostenol receives the strongest recommendation as it has reduced the 3-month rate of mortality in high-risk PAH patients also as monotherapy (2015 ESC/ERS PH guidelines). Alternative types of initial combination therapy may be considered (2015 ESC/ERS PH guidelines). Referral for lung transplantation should also be considered.

Nazzareno Galiè, Richard N. Channick, Robert P. Frantz, Ekkehard Grünig, Zhi Cheng Jing, Olga Moiseeva, Ioana R. Preston, Tomas Pulido, Zeenat Safdar, Yuichi Tamura, Vallerie V. Risk stratification and medical therapy of pulmonary arterial hypertension. *McLaughlin European Respiratory Journal* 2019 53: 1801889; DOI: 10.1183/13993003.01889-2018

CODING/BILLING INFORMATION

Note: 1) This list of codes may not be all-inclusive. 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement

HCPCS CODE	DESCRIPTION
J1325	Injection, epoprostenol, 0.5 mg
J3285	Injection, treprostinil, 1 mg
J7686	Treprostinil, inhalation solution, fda-approved final product, non-compounded, administered through dme, unit dose form, 1.74 mg
Q4074	Iloprost, inhalation solution, fda-approved final product, non-compounded, administered through dme, unit dose form, up to 20 micrograms

AVAILABLE DOSAGE FORMS:

Adcirca TABS 20MG	Sildenafil Citrate SOLN 10MG/12.5ML
Adempas TABS 0.5MG*	Sildenafil Citrate SUSR 10MG/ML
Adempas TABS 1.5MG*	Sildenafil Citrate TABS 20MG
Adempas TABS 1MG*	Tadalafil (PAH) TABS 20MG
Adempas TABS 2.5MG*	Tracleer TABS 125MG*
Adempas TABS 2MG*	Tracleer TABS 62.5MG*
Alyq TABS 20MG	Tracleer TBSO 32MG
Ambrisentan TABS 10MG	Treprostinil SOLN 100MG/20ML
Ambrisentan TABS 5MG	Treprostinil SOLN 200MG/20ML
Bosentan TABS 125MG	Treprostinil SOLN 20MG/20ML
Bosentan TABS 62.5MG	Treprostinil SOLN 50MG/20ML
Epoprostenol Sodium SOLR 0.5MG	Tyvaso SOLN 0.6MG/ML*
Epoprostenol Sodium SOLR 1.5MG	Tyvaso Refill SOLN 0.6MG/ML*
Flolan SOLR 0.5MG	Tyvaso Starter SOLN 0.6MG/ML*
Flolan SOLR 1.5MG	Uptravi SOLR 1800MCG*
Letairis TABS 10MG*	Uptravi TABS 1000MCG*
Letairis TABS 5MG*	Uptravi TABS 1200MCG*
Opsumit TABS 10MG*	Uptravi TABS 1400MCG*
Orenitram TBCR 0.125MG	Uptravi TABS 1600MCG*
Orenitram TBCR 0.25MG	Uptravi TABS 200MCG*
Orenitram TBCR 1MG	Uptravi TABS 400MCG*
Orenitram TBCR 2.5MG	Uptravi TABS 600MCG*
Orenitram TBCR 5MG	Uptravi TABS 800MCG*
Remodulin SOLN 100MG/20ML*	Uptravi TBPk 200 & 800MCG*
Remodulin SOLN 200MG/20ML*	Veletri SOLR 0.5MG
Remodulin SOLN 20MG/20ML*	Veletri SOLR 1.5MG
Remodulin SOLN 50MG/20ML*	Ventavis SOLN 10MCG/ML*
Revatio SOLN 10MG/12.5ML	Ventavis SOLN 20MCG/ML*
Revatio SUSR 10MG/ML	Tadliq SUSP 20MG/5ML
Revatio TABS 20MG	

REFERENCES

1. Adempas® tablets [prescribing information]. Whippany, NJ: Bayer; September 2021
2. Tracleer® tablets [prescribing information]. South San Francisco, CA: Actelion Pharmaceuticals, Inc; July 2022.
3. Letairis® tablets [prescribing information]. Foster City, CA: Gilead Sciences; August 2019.
4. Opsumit® tablets [prescribing information]. South San Francisco, CA: Actelion Pharmaceuticals; July 2022.
5. Flolan® injection [prescribing information]. Research Triangle Park, NC: GlaxoSmithKline; August 2021
6. Epoprostenol injection [prescribing information]. Sellersville, PA: Teva; March 2019.
7. Veletri® injection [prescribing information]. South San Francisco, CA: Actelion Pharmaceuticals; July 2022.
8. Ventavis inhalation solution [prescribing information]. South San Francisco, CA: Actelion Pharmaceuticals US, Inc; March 2022.
9. Tyvaso inhalation solution [prescribing information]. Research Triangle Park, NC: United Therapeutics Corp.; May 2022.
10. Revatio® tablets, oral suspension, and injection [prescribing information]. New York, NY: Pfizer; February 2020
11. Adcirca® tablets [prescribing information]. Indianapolis, IN: Eli Lilly (marketed by lung Biotechnology, a subsidiary of United Therapeutics Corporation); September 2020
12. Remodulin for subcutaneous or intravenous use [prescribing information]: Research Triangle Park, NC: United Therapeutics Corp; July 2021.
13. Upravi (selexipag) [package insert]. Actelion Pharmaceuticals US, Inc. San Francisco, CA. July 2022.
14. Orenitram (treprostinil) [package insert]. Research Triangle Park, NC: United Therapeutics Corp; May 2021.
15. Tadliq (tadalafil) oral suspension [prescribing information]. Farmville, NC: CMP Pharma, Inc.; June 2022.
16. Alyq (tadalafil) tablets for oral administration [prescribing information]. Jerusalem, Israel: Teva Pharmaceuticals Ind. Ltd.; September 2021.
17. McLaughlin VV, Archer SL, Badesch DB, et al; Writing committee members. ACCF/AHA 2009 Expert consensus document on pulmonary hypertension: A report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents and the American Heart Association: Developed in collaboration with the American College of Chest Physicians, American Thoracic Society, Inc., and the Pulmonary Hypertension Association. *Circulation*. 2009;119:2250-2294.
18. McGoon M, Gutterman D, Steen V, et al. Screening, early detection, and diagnosis of pulmonary arterial hypertension: ACCP evidence-based clinical practice guidelines. *CHEST*. 2004;126:14-34.
19. Badesch, Abman SH, Simonneau G, et al. Medical therapy for pulmonary arterial hypertension. Updated ACCP Evidence based clinical practice guidelines. *CHEST*. 2007;131:1917-1928.
20. Simonneau G, Gatzoulis MA, Adatia I, et al. Updated clinical classification of pulmonary hypertension. *J Am Coll Cardiol*. 2013;62(25 Suppl):D34-D41.
21. Hoeper MM, Madani MM, Nakanishi N, et al. Chronic thromboembolic pulmonary hypertension. *Lancet Respir Med*. 2014;2(7):573-582.
22. Kim NH. Group 4 pulmonary hypertension. Chronic thromboembolic pulmonary hypertension: epidemiology, pathophysiology and treatment. *Cardiol Clin*. 2016;34:435-441.
23. Simonneau G, Barst RJ, Nazzareno G, et al, for the treprostinil study group. Continuous subcutaneous infusion of treprostinil, a prostacyclin analogue, in patients with pulmonary arterial hypertension. A double-blind, randomized, placebo-controlled trial. *Am J Respir Crit Care Med*. 2002;165:800-804.
24. Barst RJ, Galie N, Naeije R, et al. Long-term outcome in pulmonary arterial hypertension patients treated with subcutaneous treprostinil. *Eur Respir J*. 2006;68:1195-1203.
25. Lang I, Gomez-Sanchez M, Kneussl M, et al. Efficacy of long-term subcutaneous treprostinil

Drug and Biologic Coverage Criteria

- sodium therapy in pulmonary hypertension. CHEST. 2006;129:1636-1643.
- Gomberg-Maitland M, Tapson VF, Benza RL, et al. Transition from intravenous epoprostenol to intravenous treprostinil in pulmonary hypertension. Am J Respir Crit Care Med. 2005;172:1586-1589.
26. Benza RL, Rayburn BK, Tallaj JA, et al. Treprostinil-based therapy in the treatment of moderate-to-severe pulmonary arterial hypertension: long-term efficacy and combination with bosentan. CHEST. 2008;134:139-145.
27. Ivy DD, Claussen L, Doran A, et al. Transition of stable pediatric patients with pulmonary arterial hypertension from intravenous epoprostenol to intravenous treprostinil. Am J Cardiol. 2007;99(5):696-698.
28. Tapson VF, Gomberg-Maitland M, McLaughlin VV, et al. Safety and efficacy of IV Treprostinil for pulmonary arterial hypertension. CHEST. 2006;129:683-688.
29. Nazzareno Galiè, Richard N. Channick, Robert P. Frantz, Ekkehard Grünig, Zhi Cheng Jing, Olga Moiseeva, Ioana R. Preston, Tomas Pulido, Zeenat Safdar, Yuichi Tamura, Vallerie V. McLaughlin. Risk stratification and medical therapy of pulmonary arterial hypertension. European Respiratory Journal 53 (1) 1801889; DOI: 10.1183/13993003.01889-2018 Published 24 January 2019
30. Adaani Frost, David Badesch, J. Simon R. Gibbs, Deepa Gopalan, Dinesh Khanna, Alessandra Manes, Ronald Oudiz, Toru Satoh, Fernando Torres, Adam Torbicki. Diagnosis of pulmonary hypertension European Respiratory Journal 53 (1) 1801904; DOI: 10.1183/13993003.01904-2018 Published 24 January 2019
31. Waxman, A., Restrepo-Jaramillo, R., Thenappan, T., Ravichandran, A., Engel, P., Bajwa, A., Allen, R., Feldman, J., Argula, R., Smith, P., Rollins, K., Deng, C., Peterson, L., Bell, H., Tapson, V., & Nathan, S. D. (2021). Inhaled Treprostinil in Pulmonary Hypertension Due to Interstitial Lung Disease. The New England journal of medicine, 384(4), 325–334.
<https://doi.org/10.1056/NEJMoa2008470>
32. Humbert, M., Kovacs, G., Hoeper, M. M., Badagliacca, R., Berger, R. M., Brida, M., Wort, S. J. (2022). 2022 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension. European Heart Journal, 43(38), 3618-3731. doi:10.1093/eurheartj/ehac237

SUMMARY OF REVIEW/REVISIONS	DATE
REVISION- Notable revisions: Products Affected Required Medical Information Continuation of Therapy Prescriber Requirements Place of Administration Drug Class FDA-Approved Uses Appendix Background Contraindications/Exclusions/Discontinuation Available Dosage Forms References	Q1 2023
Q2 2022 Established tracking in new format	Historical changes on file